Eric Borton to: Benson.Bob

```
From: Eric Borton <eric.borton@uc.edu>
  To:
 From the Lockey (1980) pleural cases, 18733, 18216, 17847, 19648,
16920
were considered bilateral.
#11377 - bilateral w/interstitial w/other asbestos
#12430 - bilateral w/interstitial
#14183 - unilateral w/interstitial
#17676 - unilateral w/interstitial
On 2/11/2011 11:18 AM, Benson.Bob@epamail.epa.gov wrote:
> Was the discrete pleural thickening found in these 4 bilateral or
unilateral? It is OK if they also had interstitial change.
> ----"Hilbert, Timothy (hilbertj)"<HILBERTJ@UCMAIL.UC.EDU>
wrote: ----
>
> To: Bob Benson/R8/USEPA/US@EPA
> From: "Hilbert, Timothy (hilbertj)"<HILBERTJ@UCMAIL.UC.EDU>
> Date: 02/11/2011 07:37AM
> Cc: "Borton, Eric (bortonek)" < BORTONEK@UCMAIL.UC.EDU>,
"Bill@mintra01.rtp.epa.gov" < Bill@mintra01.rtp.epa.gov >,
"brattin@mintra01.rtp.epa.gov" < brattin@mintra01.rtp.epa.gov >
> Subject: RE: Update bilateral pleural thickening only
>
> What would you like us to check about these 4 workers?
>
>
> ----Original Message----
> From: Benson.Bob@epamail.epa.gov
[mailto:Benson.Bob@epamail.epa.gov]
> Sent: Thursday, February 10, 2011 7:28 PM
> To: Hilbert, Timothy (hilbertj)
> Cc: Borton, Eric (bortonek); Bill@mintra01.rtp.epa.gov;
```

```
brattin@mintra01.rtp.epa.gov
> Subject: Re: Update bilateral pleural thickening only
> There may be a confusion with the terminology. I don't want
those with "Bilateral Pleural Discrete Only" but of those with
"Discete Pleural" (n = 68 with the 280 data set) who had "Bilateral
Pleural Discrete." It is OK to include those who also had
Intersitial change.
> It looks like you need to check the following workers:
> #11377
> #12430
> #14183
> #17676
> ----"Hilbert, Timothy (hilbertj)"<HILBERTJ@UCMAIL.UC.EDU>
wrote: ----
>
> To: Bob Benson/R8/USEPA/US@EPA
> From: "Hilbert, Timothy (hilbertj)"<HILBERTJ@UCMAIL.UC.EDU>
> Date: 02/10/2011 02:14PM
> Cc: "Borton, Eric (bortonek)" < BORTONEK@UCMAIL.UC.EDU>
> Subject: Update bilateral pleural thickening only
>
>
> (See attached file: Tables and Figure 02102011.xlsx)
>
> In the Rohs paper, to be counted as having bilateral pleural
changes meant at least 2 readers indicated pleural changes on both
the right and the left for any combination of discrete chest wall
pleural thickening, diaphragm pleural thickening, and/or diffuse
pleural thickening. In reviewing the 53 reported in the paper, it
was discovered that one did not meet the criteria and was only
unilateral.
> The definition utilized in the RfC is much tighter. Any workers
with asbestos exposure are excluded. Also excluded are workers who
are positive (at least 2 readers) for interstitial changes or
diffuse pleural thickening. Then to be counted as having bilateral
pleural changes meant at least 2 readers indicated only pleural
changes on both the right and the left for any combination of
discrete chest wall pleural thickening and/or diaphragm pleural
thickening.
```

```
> When we re-applied this criteria to the 252, we identified 34
subjects with only bilateral discrete pleural thickening. These
are reflected on the attached table 6. The reason for the increase
from 25 is that our previous table inadvertently excluded those
with diaphragm pleural thickening.
>
> Please let us know of any follow-up questions you may have.
> Tim
>
```